



Activity of Telavancin against Gram-Positive Cocci from Canadian Hospitals: CANWARD 2007-2010

K.A. NICHOL¹, N. LAING², B. WESHNOWESKI¹, R. VASHISHT², M.R. BAXTER², H.J. ADAM^{1,2}, D.J. HOBAN^{1,2} and G.G. ZHANEL²
Diagnostic Services of Manitoba¹ and University of Manitoba², Winnipeg, Manitoba, Canada



Kim Nichol
Health Sciences Centre
MS673 - 820 Sherbrook St.
Winnipeg, MB, Canada R3A 1R9
Phone: (204) 787-4902
Email: knichol@hsc.mb.ca

ABSTRACT

Background: Telavancin is a lipoglycopeptide with rapid bactericidal activity against a broad spectrum of Gram-positive pathogens. The purpose of this study was to assess the activity of telavancin and comparators against Gram-positive cocci obtained from Canadian hospitals as part of the ongoing national CANWARD surveillance study.

Methods: From January 2007 through December 2010, 10-15 tertiary-care centres from across Canada submitted isolates from patients attending hospital clinics, emergency rooms, medical/surgical wards, and intensive care units. Annually, each centre was asked to submit pathogens (consecutive, one per patient per infection site) from blood, respiratory specimens, urine, and wound/IV infections. 23,243 isolates were collected in total, including 10,155 Gram-positive cocci. Susceptibility testing was performed by broth microdilution using CLSI methodology for comparator agents and FDA-approved dry-form panels for telavancin.

Results: MIC₅₀ and MIC₉₀ values for telavancin, vancomycin, daptomycin and linezolid are shown below:

Organism (n)	Telavancin MIC ₅₀ /MIC ₉₀	Vancomycin MIC ₅₀ /MIC ₉₀	Daptomycin MIC ₅₀ /MIC ₉₀	Linezolid MIC ₅₀ /MIC ₉₀
MSSA (3526)	0.25 / 0.5	1 / 1	0.12 / 0.25	2 / 2
MRSA (1112)	0.25 / 0.5	1 / 1	0.25 / 0.25	2 / 2
- HA-MRSA (760)	0.25 / 0.5	1 / 1	0.12 / 0.25	2 / 2
- CA-MRSA (308)	0.25 / 0.5	1 / 1	0.25 / 0.25	2 / 2
MSSE (412)	0.25 / 0.5	1 / 2	0.12 / 0.25	0.5 / 1
MRSE (76)	0.12 / 0.25	1 / 2	0.12 / 0.25	1 / 1
<i>S. pneumoniae</i> - All (1610)	≤0.06 / ≤0.06	≤0.25 / 0.25	≤0.06 / 0.12	0.5 / 1
- PenS (1301)	≤0.03 / ≤0.06	≤0.25 / 0.25	≤0.06 / ≤0.06	0.5 / 1
- PenI (224)	≤0.06 / ≤0.06	≤0.25 / 0.5	0.06 / 0.12	1 / 1
- PenR (72)	≤0.03 / ≤0.06	≤0.25 / 0.5	0.06 / 0.12	0.5 / 1

MSSA, methicillin-susceptible *S. aureus*; MRSA, methicillin-resistant *S. aureus*; HA-MRSA, healthcare-associated MRSA; CA-MRSA, community-associated MRSA; MSSE, methicillin-susceptible *S. epidermidis*; MRSE, methicillin-resistant *S. epidermidis*.

Conclusions: Telavancin is more active in vitro than vancomycin and linezolid and has comparable activity to daptomycin against MSSA, MRSA (including CA-MRSA and HA-MRSA), MSSE, MRSE and *S. pneumoniae*.

BACKGROUND

Antibiotic resistance among Gram-positive pathogens such as *Streptococcus pneumoniae*, *Staphylococcus aureus* and *Staphylococcus epidermidis* is a growing concern. The global escalation in both community- and healthcare-acquired antibiotic-resistant organisms is threatening our ability to effectively treat patients by significantly limiting the therapeutic options available to clinicians and increasing the risk of treatment failures. Management of infections caused by these difficult-to-treat pathogens is often complicated further by the fact that many of these strains are multidrug-resistant. These observations underscore the need for continued surveillance, more judicious antibiotic prescribing practices and new treatment alternatives.

Telavancin is a semisynthetic lipoglycopeptide with a dual mechanism of action against a broad spectrum of clinically relevant Gram-positive bacteria, including both susceptible and multidrug-resistant staphylococci and streptococci (1-4). The rapid bactericidal activity of telavancin is derived from its ability to inhibit synthesis of the bacterial cell wall as well as to disrupt bacterial membrane integrity and increase cell membrane permeability (1-4).

PURPOSE

The purpose of this study was to assess the activity of telavancin and comparators against Gram-positive cocci obtained from Canadian hospitals in 2007-2010 as part of the ongoing national CANWARD surveillance study.

MATERIALS & METHODS

CANWARD Study Design

Between January 2007 and December 2010, 23,243 clinical isolates, including 10,155 Gram-positive cocci, were collected as part of the ongoing CANWARD study assessing pathogen prevalence and antibiotic resistance in Canadian hospitals. Isolates were received from tertiary-care medical centres (12 in 2007, 10 in 2008, 15 in 2009, 14 in 2010) that were geographically distributed in a population-based fashion in eight of the ten Canadian provinces. Annually, each study site was asked to submit clinical isolates (consecutive, one per patient per infection site) from inpatients and outpatients with respiratory, urine, wound and bloodstream infections. Isolates were collected from patients attending hospital clinics, emergency rooms, medical/surgical wards and intensive care units. All organisms were identified by the submitting centre and were deemed clinically significant using local site criteria. Isolates were shipped on Amies semi-solid transport media to the coordinating laboratory (Health Sciences Centre, Winnipeg, Canada) where they were subcultured onto appropriate media and stocked in skim milk at -80°C.

Antimicrobial Susceptibility Testing

Following two subcultures from frozen stock, the *in vitro* activities of telavancin and comparator agents, including cefazolin, ceftriaxone, clarithromycin, clindamycin, ciprofloxacin, daptomycin, doxycycline, ertapenem, levofloxacin, linezolid, meropenem, moxifloxacin, penicillin, piperacillin-tazobactam, tigecycline, trimethoprim-sulfamethoxazole and vancomycin, were determined by broth microdilution in accordance with Clinical and Laboratory Standards Institute (CLSI) guidelines (5-6). Antimicrobial minimum inhibitory concentrations (MICs) were determined using 96-well custom designed microtitre panels for comparator agents and FDA-approved dry-form panels for telavancin. Quality control was performed using the following ATCC organisms: *S. pneumoniae* ATCC 49619 and *S. aureus* ATCC 29213. MIC interpretive standards were defined according to CLSI breakpoints (7). The following interpretive breakpoints (FDA) were used with *S. aureus*: telavancin susceptible, ≤1 µg/ml; tigecycline susceptible, ≤0.5 µg/ml. With *S. pneumoniae*, an interpretive breakpoint (FDA) of ≤0.06 µg/ml was used for tigecycline susceptible. Tetracycline breakpoints were used to interpret *S. pneumoniae* doxycycline MIC values.

ACKNOWLEDGMENTS

Financial support for the CANWARD study was provided in part by Astellas Pharma Inc.

The authors would like to thank the participating centres, investigators and laboratory site staff for their continued support:

Dr. D. Roscoe – Vancouver Hospital, Vancouver	Dr. M. Desjardins – The Ottawa Hospital, Ottawa
Dr. R. Rennie – University of Alberta Hospital, Edmonton	Dr. M. Laverdière – Hôpital Maisonneuve-Rosemont, Montreal
Dr. J. Blondeau – Royal University Hospital, Saskatoon	Dr. V. Loo – Montreal General Hospital, Montreal
Drs. D. Hoban/G. Zhanel – Health Sciences Centre, Winnipeg	Dr. V. Loo – Royal Victoria Hospital, Montreal
Dr. Z. Hussain – London Health Sciences Centre, London	Dr. M. Goyette – CHRTR Pavillon Ste. Marie, Trois-Rivières
Dr. S. Poutanen – Mount Sinai Hospital, Toronto	Dr. M. Kuhn – South East Regional Health Authority, Moncton
Dr. L. Matukas – St. Michael's Hospital, Toronto	Dr. R. Davidson – Queen Elizabeth II HSC, Halifax
Dr. F. Chan – Children's Hospital of Eastern Ontario, Ottawa	

Table 1. Activity of telavancin and comparators against Gram-positive cocci from CANWARD 2007-2010

Organism (n), Antibiotic	% of Isolates per Category			MIC ₅₀	MIC ₉₀	Range Min	Range Max
	S	I	R				
S. aureus							
MSSA (3526)							
Telavancin ^a	100			0.25	0.5	≤ 0.06	1
Vancomycin	100			1	1	≤ 0.25	2
Cefazolin	100			≤ 0.5	1	≤ 0.5	32
Clarithromycin	74.5	0.3	25.2	0.25	> 16	≤ 0.25	> 16
Clindamycin	91.9	0.5	7.7	≤ 0.25	≤ 0.25	≤ 0.25	> 8
Daptomycin	100			0.12	0.25	≤ 0.06	1
Levofloxacin	90.1	0.3	9.6	0.25	1	≤ 0.06	> 32
Linezolid	100			2	2	≤ 0.12	4
Meropenem	100			0.12	0.12	≤ 0.12	4
Moxifloxacin	90.0	0.7	9.3	0.06	1	≤ 0.06	> 16
Pip-Tazo	99.9		0.1	≤ 1	≤ 1	≤ 1	32
Tigecycline ^a	99.9			0.25	0.25	≤ 0.03	1
TMP/SMX	99.4		0.6	≤ 0.12	≤ 0.12	≤ 0.12	> 8
MRSA (1112)							
Telavancin ^a	100			0.25	0.5	0.12	1
Vancomycin	99.9	0.1		1	1	≤ 0.25	4
Cefazolin			100.0 ^b	64	> 128	1	> 128
Clarithromycin	11.8	0.1	88.1	> 16	> 16	≤ 0.25	> 16
Clindamycin	46.3	0.1	53.6	> 8	> 8	≤ 0.25	> 8
Daptomycin	100			0.25	0.25	0.06	1
Levofloxacin	14.1			> 32	> 32	0.12	> 32
Linezolid	100			2	2	≤ 0.12	4
Meropenem				8	32	0.12	> 32
Moxifloxacin	14.5	1.6	83.9	8	> 16	≤ 0.06	> 16
Pip-Tazo	100.0 ^b			64	128	≤ 1	512
Tigecycline ^a	99.8			0.25	0.5	0.06	1
TMP/SMX	91.4		8.6	≤ 0.12	0.25	≤ 0.12	> 8
CA-MRSA (308)							
Telavancin ^a	100			0.25	0.5	0.12	1
Vancomycin	100			1	1	0.5	2
Cefazolin			100.0 ^b	16	64	1	> 128
Clarithromycin	24.4	0.3		75.3	> 16	> 16	≤ 0.25
Clindamycin	86.7			13.3	≤ 0.25	> 8	≤ 0.25
Daptomycin	100			0.25	0.25	0.12	1
Levofloxacin	39.9			60.1	4	8	0.12
Linezolid	100			2	2	1	4
Meropenem			100.0 ^b	2	4	0.12	16
Moxifloxacin	37.0	5.5		57.5	2	2	≤ 0.06
Pip-Tazo	100		100.0 ^b	16	32	2	64
Tigecycline ^a	100			0.25	0.25	0.06	0.5
TMP/SMX	100			≤ 0.12	≤ 0.12	≤ 0.12	2
HA-MRSA (760)							
Telavancin ^a	100			0.25	0.5	0.12	1
Vancomycin	99.9	0.1		1	1	≤ 0.25	4
Cefazolin			100.0 ^b	128	> 128	1	> 128
Clarithromycin	4.5			95.5	> 16	> 16	≤ 0.25
Clindamycin	28.8	0.1		71.1	> 8	> 8	≤ 0.25
Daptomycin	100			0.12	0.25	0.06	1
Levofloxacin	2.9			97.1	> 32	> 32	0.12
Linezolid	100			2	2	≤ 0.12	4
Meropenem			100.0 ^b	16	> 32	0.25	> 32
Moxifloxacin	2.9	0.1		97	> 16	≤ 0.06	> 16
Pip-Tazo	99.7		100.0 ^b	64	128	≤ 1	512
Tigecycline ^a	99.7			0.25	0.5	0.12	1
TMP/SMX	87.5		12.5	≤ 0.12	8	≤ 0.12	> 8

^a Interpretive breakpoints defined by FDA.

^b Based on oxacillin susceptibility.

S, susceptible; I, intermediate; R, resistant.

CA-MRSA, community-associated methicillin-resistant *S. aureus*.
HA-MRSA, healthcare-associated methicillin-resistant *S. aureus*.

Table 2. MIC distribution of telavancin against Gram-positive cocci from CANWARD 2007-2010

Organism (no. tested)	Number (cumulative percentage) at each MIC				
	≤ 0.06	0.12	0.25	0.5	1
Methicillin-susceptible <i>S. aureus</i> (2801)	15 (0.5)	1000 (36.2)	1115 (76.0)	573 (96.5)	98 (100)
Methicillin-resistant <i>S. aureus</i> (852)		246 (28.9)	343 (69.1)	228 (95.9)	35 (100)
Community-associated MRSA (258)		80 (31.0)	105 (71.7)	64 (96.5)	9 (100)
Healthcare-associated MRSA (562)		156 (27.8)	223 (67.4)	157 (95.4)	26 (100)
Methicillin-susceptible <i>S. epidermidis</i> (345)	36 (10.4)	130 (48.1)	122 (83.5)	50 (98.0)	7 (100)
Methicillin-resistant <i>S. epidermidis</i> (65)	3 (4.6)	30 (50.8)	26 (90.8)	3 (95.4)	3 (100)
<i>S. pneumoniae</i> (1610)	1608 (99.9)	2 (100)			
Methicillin-susceptible <i>S. pneumoniae</i> (1298)	1296 (99.8)	2 (100)			
Penicillin-intermediate <i>S. pneumoniae</i> (224)	224 (100)				
Penicillin-resistant <i>S. pneumoniae</i> (72)	72 (100)				

MIC, minimum inhibitory concentration (µg/ml).

Organism (n), Antibiotic	% of Isolates per Category			MIC ₅₀	MIC ₉₀	Range Min	Range Max
	S	I	R				
<i>S. pneumoniae</i>							
All (1610)							
Telavancin	No BP			≤ 0.06	≤ 0.06	≤ 0.06	0.12
Vancomycin	100			≤ 0.25	0.25	≤ 0.25	1
Ceftriaxone	99.4	0.4	0.2	≤ 0.12	≤ 0.12	≤ 0.12	4
Clarithromycin	80.7	4.1	15.2	≤ 0.03	2	≤ 0.03	> 32
Clindamycin	93.6	0.6	5.8	≤ 0.12	≤ 0.12	≤ 0.12	> 8
Doxycycline ^c	93.2	3.1	3.7	≤ 0.25	1	≤ 0.25	> 16
Levofloxacin	99.1	0.1	0.8	0.5	1	≤ 0.06	32
Linezolid	100			0.5	1	≤ 0.12	2
Meropenem	96.1	2.5	1.4	≤ 0.06	≤ 0.06	≤ 0.06	2
Penicillin	81.5	14.0	4.5	≤ 0.03	0.25	≤ 0.03	> 8
Tigecycline ^a	99.7			0.03	0.06	≤ 0.03	0.25
TMP/SMX	84.8	6.6	8.6	≤ 0.12	2	≤ 0.12	> 8
PSSP (1301)							
Telavancin	No BP			≤ 0.03	≤ 0.06	≤ 0.03	0.12
Vancomycin	100			≤ 0.25	0.25	≤ 0.12	1
Ceftriaxone	100			≤ 0.06	≤ 0.12	≤ 0.06	0.25
Clarithromycin	88.5	3.2	8.4	≤ 0.03	0.5	≤ 0.03	> 32
Clindamycin	98.0	0.4	1.6	≤ 0.12	≤ 0.12	≤ 0.12	> 16
Doxycycline ^c	97.4	1.5	1.2	≤ 0.25	≤ 0.25	≤ 0.25	> 16
Levofloxacin	99.3	0.2	0.5	0.5	1	≤ 0.06	32
Linezolid	100			0.5	1	≤ 0.12	2
Meropenem	99.9	0.1		≤ 0.06	≤ 0.06	≤ 0.06	0.5
Penicillin	100.0			≤ 0.03	0.06	≤ 0.03	0.06
Tigecycline ^a	99.9			0.03	0.06	≤ 0.015	0.25
TMP/SMX	91.4	5.7	2.9	≤ 0.12	0.5	≤ 0.12	> 8
PISP (224)							
Telavancin	No BP			≤ 0.06	≤ 0.06	≤ 0.06	0.06
Vancomycin	100			≤ 0.25	0.5	≤ 0.12	1
Ceftriaxone	99.1	0.5	0.5	≤ 0.12	0.25	≤ 0.12	4
Clarithromycin	49.3	9.9	40.8	0.5	> 32	≤ 0.03	> 32
Clindamycin	79.8	1.4	18.8	≤ 0.12	> 8	≤ 0.12	> 8
Doxycycline ^c	75.3	6.3	18.4	≤ 0.25	16	≤ 0.25	> 16
Levofloxacin	97.8		2.2	1	1	0.25	16
Linezolid	100			1	1	≤ 0.12	2
Meropenem	95.1	4.0	0.9	≤ 0.06	0.25	≤ 0.06	1
Penicillin	100			0.12	1	0.12	1
Tigecycline ^a	99.1			0.03	0.06	≤ 0.03	0.12
TMP/SMX	67	11.6	21.4	0.25	8	≤ 0.12	> 8
PRSP (72)							
Telavancin	No BP			≤ 0.03	≤ 0.06	≤ 0.03	0.06
Vancomycin	100			≤ 0.25	0.5	≤ 0.25	1
Ceftriaxone	88.9	8.3	2.8	0.5	2	≤ 0.06	4
Clarithromycin	36.1	2.8	61.1	4	> 32	≤ 0.03	> 32
Clindamycin	56.9	2.8	40.3	≤ 0.12	> 64	≤ 0.12	> 64
Doxycycline ^c	73.6	22.2	1	4	5	≤ 0.25	> 16
Levofloxacin	98.6		1.4	1	1	0.25	16
Linezolid	100			0.5	1	0.25	2
Meropenem	30.6	40.3	29.2	0.5	1	≤ 0.06	2
Penicillin	100			2	4	> 2	> 8
Tigecycline ^a	98.6			0.03	0.06	≤ 0.015	0.12
TMP/SMX	18.1	6.9	75.0	4	> 8	≤ 0.12	> 8