

Activity of Doripenem (DOR) Against Molecularly Characterized AmpC and ESBL Producing *Escherichia coli* (EC) Including Strains with Reduced Susceptibility to Carbapenems (CRS) from Canadian Hospitals: CANWARD 2007-2010

F. TAILOR¹, P. J. SIMNER¹, L. MATASEJE², M. R. MULVEY², P. LAGACÉ-WIENS³, M. DECORBY¹, B. WESHNOWESKI⁴, R. VASHISHT¹, N. LAING¹, D. J. HOBAN⁴ and G. G. ZHANEL¹

¹University of Manitoba, ²National Microbiology Laboratory, ³St-Boniface Hospital and ⁴Health Sciences Centre, Winnipeg, Canada

ABSTRACT

Background: DOR is a recently approved carbapenem. The purpose of this study was to determine the activity of DOR against CRS-EC, AmpC-EC and ESBL-EC.

Methods: From 2007-2010 inclusive, a total of 4806 EC were collected from medical centres across Canada as part of a national surveillance program assessing antimicrobial resistance in Canadian hospitals. Susceptibility testing was performed using CLSI methods. PCR and sequencing of β -lactamase genes was used to determine AmpC and ESBL genotypes. Isolates with an ertapenem MIC ≥ 0.12 μ g/ml were defined as having reduced susceptibility to carbapenems (CRS) and were tested for KPC by PCR. SDS-PAGE and sequencing assessed OmpF and OmpC among CRS isolates. PFGE was used to determine genetic relatedness.

Results: Activity of DOR and comparators:

Drug	MIC ₅₀ /MIC ₉₀ (μ g/ml)			
	Total EC (n=4807)	AmpC-EC (n=97)	ESBL-EC (n=186)	CRS-EC (n=57)
AMC ^a	4/8 ^c	16/>32	8/16	16/>32
Cefepime	$\leq 1/\leq 1^d$	$\leq 1/\leq 1^f$	16/>32 ^h	16/>32 ⁱ
Ceftazidime	$\leq 0.5/1^e$	2/>32 ^g	16/>32 ⁱ	>32/>32
Ceftobiprole	$\leq 1/\leq 1$	$\leq 1/\leq 1$	32/>32	18/>32
Ceftriaxone	$\leq 1/\leq 1$	8/64	>64/>64	>64/>64
Cefoxitin	4/8 ^c	>32/>32	8/32	32/>32
Doripenem	$\leq 0.12/\leq 0.12$	$\leq 0.12/\leq 0.12$	$\leq 0.12/\leq 0.12$	$\leq 0.12/\leq 0.12$
Ertapenem	$\leq 0.06/\leq 0.06^e$	$\leq 0.06/0.25$	$\leq 0.06/0.25$	0.25/1
Meropenem	$\leq 0.12/\leq 0.12$	$\leq 0.12/\leq 0.12$	$\leq 0.12/\leq 0.12$	$\leq 0.12/\leq 0.12$
TZP ^b	2/4	4/32	4/16	8/64

^a: AMC, Amoxicillin/Clavulanic Acid; ^b: TZP, Piperacillin Tazobactam; ^c: n=3805; ^d: n=3788;

^e: n=3247; ^f: n=70; ^g: n=93; ^h: n=155; ⁱ: n=134; ^j: n=46

Conclusions: DOR displayed potent *in vitro* activity against CRS-EC, AmpC-EC and ESBL-EC isolates and displayed comparable activity to meropenem. DOR represents a potential option for infections caused by CRS-EC, AmpC-EC and ESBL-EC isolates.

ACKNOWLEDGMENTS

The CANWARD study was supported in part by Abbott, Astellas, AstraZeneca, Cerexa/Forest, Merck, Pfizer, Sepracor, The Medicines Company and Affinium. The authors would also like to thank the participating centres, investigators, and laboratory staff for their continued support and The Canadian Antimicrobial Resistance Alliance (CARA).

REFERENCES

- Toye, B., et al. *Can. Med. Assoc. J.* 2009. **180**:1225-1226.
- Paterson, D., et al. *Clin. Infect. Dis.* 2009. **49**:291-298.
- Nordmann, P., et al. *Lancet Infect. Dis.* 2009. **9**:228-238.
- Queenan, A. M., et al. *Clin. Microbiol. Rev.* 2007. **20**:440-458.
- Falagas, M. E., et al. *Clin. Infect. Dis.* 2005. **40**:1333-1341.
- Steward, C. D., et al. *J. Clin. Microbiol.* 2001. **39**:2864-2872.
- Perez-Perez, F. J., et al. *J. Clin. Microbiol.* 2002. **40**:2153-2162.
- Caroff, N., et al. *FEMS Microbiol.* 1999. **173**:459-465.
- Mulvey, M. R., et al. *Antimicrob. Agents Chemother.* 2005. **49**:358-365.
- Hernández-Allés, S., et al. *Microbiol.* 1999. **143**:673-679.
- Doumith, M., et al. *J. Antimicrob. Chemother.* 2009. **63**:659-667.

BACKGROUND

In the last two decades, carbapenems have been considered the last line of defence against multi-drug resistant (MDR-defined as concomitant resistance to ≥ 3 different antimicrobial classes) infections caused by *Enterobacteriaceae* (1). From this family, *Escherichia coli* (EC) remains the leading pathogen causing infections in Canada (<http://www.can-r.ca>). EC is associated with many infections including urinary tract infections, bacteremia, pneumonia, meningitis, and diarrheal diseases. Traditional treatment includes the use of β -lactam antibiotics (penicillins, cephalosporins) with the main concern being the development of resistance towards these agents with production of extended-spectrum β -lactamases (ESBLs) and/or AmpC β -lactamases by these organisms. Doripenem, a recently approved carbapenem, has been shown to have intrinsic stability to most β -lactamases including ESBLs and AmpCs. Its antimicrobial spectrum most closely resembles meropenem (MER) and imipenem (IPM) than ertapenem (ETP) (2). With the increase in use of carbapenems, the greatest concern is the development of carbapenem resistance or reduced susceptibility (defined as an MIC of ETP ≥ 0.12 μ g/ml). Carbapenem resistance can arise through the production of carbapenemases which include the *Klebsiella pneumoniae* carbapenemases (KPCs), as well as ESBL and/or AmpC production coupled with porin modification (3). Genes encoding KPCs are carried on plasmids and have the ability to hydrolyze virtually all β -lactam antibiotics including carbapenems (3, 4). Plasmids may also carry other resistant determinants such as resistance to aminoglycosides, fluoroquinolones, tetracyclines, and trimethoprim sulfamethoxazole conferring a MDR profile. For this reason, treatment can be very problematic and limited to only few agents such as tigecycline and colistin (1, 5). The purpose of this study was to determine the DOR susceptibilities of CRS-EC, ESBL-EC and AmpC-EC versus comparator antimicrobial agents.

MATERIALS & METHODS

Bacterial Isolates: Isolates were selected from the Canadian Ward (CANWARD) surveillance study which is a national study based at Health Sciences Centre (HSC) in Winnipeg, MB, Canada. From January 2007 through November 2010, participating Canadian hospitals submitted pathogens from patients attending hospital clinics, emergency rooms, medical and surgical wards, and intensive care units. Annually, each centre was asked to submit pathogens (consecutive, one per patient/infection site) from blood, respiratory, urine, and wound/IV infections. All isolates were identified at the originating centre using routine procedures performed at each site. A total of 7718, 5282, 5375, and 4868 isolates were collected for 2007, 2008, 2009, and 2010, respectively. Of these, 4807 were *E. coli* (EC).

Susceptibility Testing: The *in vitro* activity of various antimicrobials were determined (CRS isolates tested in duplicate to confirm ETP results) by broth microdilution in accordance with CLSI guidelines (CLSI M07-A8, Vol.29 No.2). The minimum inhibitory concentration (MIC) was determined using custom designed 96-well microtitre plates. 2010 CLSI breakpoints were used for interpretation excluding the ones mentioned following. Food and Drug Administration interpretation breakpoints were used for tigecycline (S: ≤ 2 μ g/ml, I: 4 μ g/ml, R: ≥ 8 μ g/ml) and colistin (S: ≤ 2 μ g/ml, R: ≥ 4 μ g/ml). Health Canada interpretation breakpoints were used for ceftobiprole (S: ≤ 1 μ g/ml, I: 2 μ g/ml, R: ≥ 4 μ g/ml). Any EC with a ceftriaxone or ceftazidime MIC ≥ 1 μ g/ml were further tested using disk diffusion methods described by CLSI to confirm ESBL phenotype (CLSI M100-S21, Vol.31 No.1.). Any EC with a ceftazidime MIC ≥ 32 μ g/ml was identified as a putative AmpC-producer. Any EC with an ertapenem MIC ≥ 0.12 μ g/ml was defined as carbapenem reduced susceptible.

Characterization of ESBLs and AmpCs: Genotypic characterization of ESBLs was performed by PCR and sequencing of *bla*_{SHV}, *bla*_{TEM}, *bla*_{CTX-M}, and *bla*_{OXA} genes as previously described (6). All isolates with a MIC ≥ 32 μ g/ml were screened for genes encoding EBC (ACT-1/MIR-1-related genes), DHA (DHA-related genes), FOX (FOX-related genes) and CIT (CMY-2-related genes) phylogenetic groups of AmpC acquired enzymes using a previously described multiplex PCR (7). A BLAST search of the DNA sequence was conducted to determine specific genotypes.

PCR for the amplification of the ampC promoter region was performed as previously described on plasmidic AmpC negative isolates (8).

Characterization of CRS-EC: Phenotypic characterization of CRS-EC was performed using Modified Hodge Test (MHT). Genotypic characterization was performed by KPC PCR.

Molecular Subtyping by PFGE: Genetic relationships of the ESBL-, AmpC-producing and CRS-EC were assessed by pulsed-field gel electrophoresis (PFGE) following digestion with *Xba*I as previously described (9).

Outer Membrane Porin (OMP) Analysis of CRS-EC: Cells grown in nutrient broth overnight at 37°C were sonicated and OMPs were analyzed using SDS-PAGE and sequencing of OmpF/C of EC (10, 11).

RESULTS

Table 1. Antimicrobial profiles among total-EC, AmpC-EC, ESBL-EC, and CRS-EC from CANWARD 2007-2010

Drug	Total EC (n=4807) ^a					AmpC-EC (n=97) ^b					ESBL-EC (n=186) ^c					CRS-EC (n=57) ^d									
	MIC 50/90 μ g/ml	% S	I	R	% S I R	MIC 50/90 μ g/ml	% S	I	R	% S I R	MIC 50/90 μ g/ml	% S	I	R	% S I R	MIC 50/90 μ g/ml	% S	I	R	% S I R					
Cefazolin	2	16	36	33.8	30.1	>128	>128	1	0	99	>128	>128	0	0	100	>128	>128	0	0	100	>128	>128	0	0	100
Ceftazidime	≤ 0.5	1	95.6	0.6	3.8	2	>32	67.7	1.1	31.2	16	>32	36.6	9	54.5	>32	>32	15.8	3.5	80.7	>32	>32	15.8	3.5	80.7
Ceftriaxone	≤ 1	≤ 1	91.1	0.5	8.5	8	>64	42.3	1	56.7	>64	>64	1.6	2.2	96.2	>64	>64	10.5	0	89.5	>64	>64	10.5	0	89.5
Cefepime	≤ 1	≤ 1	96.6	1.9	1.5	0.25	1	97.1	1.4	1.4	16	>32	49	27.7	23.2	16	>32	48.9	6.7	44.4	>32	>32	48.9	6.7	44.4
Ceftobiprole	≤ 1	≤ 1	95.9	0.2	3.9	0.12	1	91.8	3.1	5.2	32	>32	3.2	3.8	93	8	>32	38.6	5.3	56.1	>32	>32	38.6	5.3	56.1
Cefoxitin	4	8	91.9	3.9	4.1	>32	>32	0	0	100	8	32	81.2	8.6	10.2	32	>32	15.8	19.3	64.9	>32	>32	15.8	19.3	64.9
Doripenem	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0
Ertapenem	≤ 0.06	≤ 0.06	99.5	0.2	0.3	≤ 0.06	0.25	93.8	2.1	4.1	≤ 0.06	0.25	94.1	3.2	2.7	0.25	1	70.2	12.3	17.5	≤ 0.06	≤ 0.06	99.5	0.2	0.3
Meropenem	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0	≤ 0.12	≤ 0.12	100	0	0
Amoxicillin/Clav.	4	8	90.1	6.9	3	16	>32	34	24.7	41.2	8	16	67.7	27.4	4.8	16	>32	35.1	29.8	35.1	>32	>32	35.1	29.8	35.1
Piperacillin/Tazo.	2	4	97.6	1.3	1.1	4	32	89.7	7.2	3.1	4	16	91.9	5.9	1.6	8	64	68.4	22.8	8.8	>32	>32	68.4	22.8	8.8
Ciprofloxacin	≤ 0.06	>16	78.6	0.3	21.1	0.25	>16	56.7	2.1	41.2	>16	>16	12.4	0.5	87.1	>16	>16	28.1	1.8	70.2	>16	>16	28.1	1.8	70.2
Levofloxacin	≤ 0.06	>16	78.8	0.7	20.6	2	>32	50	2.9	47.1	16	32	12.9	1.3	85.8	16	>32	24.2	0	75.6	>32	>32	24.2	0	75.6
Trimethoprim/Sulf.	≤ 0.12	>8	73.7	-	26.3	0.25	>8	66	-	34	>8	>8	28	-	72	1	>8	54.4	-	45.6	>8	>8	54.4	-	45.6
Tigecycline	0.5	1	99.9	0.1	0	0.5	1	99	1	0	0.5	1	99.5	0.5	0	0.5	1	98.2	1.8	0	>8	>8	98.2	1.8	0
Amikacin	≤ 2	4	99.6	0.3	0.2	2	4	97.9	0	2.1	4	16	95.7	3.8	0.5	2	16	91.2	5.3	3.5	>32	>32	91.2	5.3	3.5
Gentamicin	≤ 0.5	2	90.9	0.3	8.8	0.5	32	81.4	0	18.6	16	>32	50.5	0.5	48.9	2	>32	59.6	0	40.4	>32	>32	59.6	0	40.4
Colistin	0.5	0.5	99.4	0	0.6	0.25	0.5	100	-	0	0.5	1	98.9	-	1.1	0.5	1	98.2	-	1.8	>32	>32	98.2	-	1.8

^aceftazidime, n=3247; cefepime, n=3788; cefoxitin, n=3805; ertapenem, n=3803 amoxicillin/clav., n=3805; levofloxacin, n=3788; colistin, n=3805

^bceftazidime, n=93; cefepime, n=70; levofloxacin, n=70

^cceftazidime, n=134; cefepime, n=155; levofloxacin, n=155

^dcefepime, n=45; levofloxacin, n=45

Figure 1. Genetic relationship among CRS-EC from CANWARD 2007-2010




Table 2. Porin profiles of CRS-EC from CANWARD 2007-2010

Isolate #	Molecular mechanisms	MIC (μ g/ml) ^a								OmpF (genetic lesion) ^b	OmpC (genetic lesion) ^b	
		ETP	MEM	DOR	CRO	CAZ	FEP	FOX	AMC			TZP
76819	CTX-M-15, OXA-1	0.25	≤ 0.06	≤ 0.06	>64	32	32	4	16	4	A	ins \rightarrow PS
81687	CTX-M-15, OXA-1	0.25	0.12	0.12	>64	>32	>32	16	8	128	NEG	TTGGG del \rightarrow PS
82395	CTX-M-14, TEM-1	0.12	≤ 0.06	≤ 0.06	64	2	>32	32	4	4	GAAC	ins \rightarrow PS
84251	CMY-2	0.12	≤ 0.06	≤ 0.06	16	32	≤ 0.25	>32	8	8	NS mut	NEG
85332	CTX-M-14, TEM-1	0.12	≤ 0.06	≤ 0.06	32	2	4	32	16	8	GAAC	ins \rightarrow PS
86230	Unknown	0.12	≤ 0.06	≤ 0.06	≤ 0.25	1	≤ 0.25	16	16	32	NEG	A \rightarrow G (RRPM)
90484	-18 (G to A), -1 (C to A), +58 (C to T)	0.12	≤ 0.06	≤ 0.06	0.5	8	-	>32	>32	≤ 1	NEG	A \rightarrow G (RRPM)

Figure 2. Resistance mechanisms among CRS-EC isolates (CANWARD 2007-2010)

Outer Membrane Porin Profiles

E. coli

Approximately 30-35% of isolates were shown to have at least 1 of OmpF or OmpC missing. Sequencing showed insertions, deletions, and nonsense mutations causing premature termination.

